

Cost effectiveness of targeted HIV prevention interventions for female sex workers in India

Shankar Prinja,¹ Pankaj Bahuguna,¹ Shalini Rudra,² Indrani Gupta,² Manmeet Kaur,¹ S M Mehendale,³ Susmita Chatterjee,⁴ Samiran Panda,⁴ Rajesh Kumar¹

► Additional materials are published online only. To view these files please visit the journal online (<http://sti.bmj.com>).

¹School of Public Health, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

²Institute of Economic Growth, Delhi University, New Delhi, India

³National Institute of Epidemiology, Chennai, India

⁴National Institute of Cholera and Enteric Diseases (NICED), Kolkata, India

Correspondence to

Dr Shankar Prinja, School of Public Health, Post Graduate Institute of Medical Education and Research, Chandigarh 160012, India; shankarprinja@gmail.com

Accepted 13 February 2011

ABSTRACT

Objective To ascertain the cost effectiveness of targeted interventions for female sex workers (FSW) under the National AIDS Control Programme in India.

Methods A compartmental mathematical Markov state model was used over a 20-year time horizon (1995–2015) to estimate the cost effectiveness of FSW targeted interventions, with a health system perspective. The incremental costs and effects of FSW targeted interventions were compared against a baseline scenario of mass media for the general population alone. The incremental cost-effectiveness ratio was computed at a 3% discount rate using HIV infections averted and disability-adjusted life-years (DALY) as benefit measures. It was assumed that the transmission of the HIV virus moves from a high-risk group (FSW) to the client population and finally to the general population (partners of clients).

Result Targeted interventions for FSW result in a reduction of 47% (1.6 million) prevalent and 36% (2.7 million) cumulative HIV cases, respectively, in 2015. Adult HIV prevalence in India, with and without (mass media only) FSW interventions, would be 0.25% and 0.48% in 2015. Indian government and development partners spend an average US\$104 (INR4680) per HIV infection averted and US\$10.7 (INR483) per DALY averted. Discounting at 3%, FSW targeted interventions cost US\$105.5 (INR4748) and US\$10.9 (INR490) per HIV case and DALY averted, respectively.

Conclusion At the current gross domestic product in India, targeted intervention is a cost-effective strategy for HIV prevention in India.

India initiated the National AIDS Control Programme in 1992, which is currently in the third phase of implementation. The total budgetary outlay of the programme has increased from INR800 million to INR115 830 million from the first to the third phase.¹ Prevention has been the mainstay of the programme. Gradually, the major strategy for prevention has shifted from creating awareness through mass media towards a behaviour change communication and providing an enabling environment through peer-led targeted interventions. These targeted interventions have focused on high-risk groups of female sex workers (FSW), men who have sex with men (MSM) and injecting drug users (IDU). The main service targeted interventions for FSW include peer-led counselling for behaviour change towards safer sexual practices, condom promotion, quarterly referral for health check-up and sexually transmitted infection (STI) treatment, and referral and support for accessing antiretroviral therapy (ART)

after self-disclosure of HIV-positive status by the FSW to an outreach peer worker. Currently, there are more than 1500 targeted intervention projects, of which more than 1200 are funded by the National AIDS Control Organization (NACO).²

Case studies from India have shown the effectiveness of targeted interventions for FSW in reducing risky sexual behaviour.^{3–5} Other studies have used mathematical modelling to relate improvements in behaviour, with a declining trend in HIV incidence and prevalence.^{6–7} A significantly declining trend in HIV prevalence among FSW and the general population was observed in districts that had a high intensity of preventive programmes compared with low intensity districts in four south Indian high HIV prevalence states.²

Other studies conducted in developing and developed countries have also shown that preventive strategies promote safer sexual behaviours⁸ and are also cost effective.^{9–10} State-level analyses in India have recently reported targeted interventions to be cost effective in Gujarat state.¹¹ More research has also been carried out to estimate the cost of scale-up of targeted interventions in the former state.¹² However, no national-level study has been conducted so far, to inform resource allocation for the country's preventive interventions under the National AIDS Control Programme. The present study was thus conducted to ascertain the incremental cost effectiveness of targeted interventions in India. As female sex work is the major driver of India's HIV epidemic,^{13–15} a focus on FSW targeted interventions was considered appropriate for the present study.

METHODS

General model description

A dynamic Markov model was parameterised on an MS Excel spreadsheet to estimate the incremental cost effectiveness of FSW targeted interventions. Incremental costs and the effectiveness of FSW interventions were estimated against a comparator scenario of mass media interventions alone. A time horizon of 20 years was considered, starting from the base year of 1995. The present study takes a health system perspective. Benefit was measured in terms of HIV infections averted and disability-adjusted life-years (DALY) averted. Both costs and benefits were discounted at 3% to account for time preference of cost and utility.¹⁶

Demographic and epidemiological assumptions

We assume the transmission of HIV virus from a high-risk group (FSW) to the client population

and finally to the general population (partners of clients) (figure 1). We also include reverse transmission of HIV from clients to FSW, but assume no transmission of HIV from the general population to the client population.

We used census growth rates and behavioural parameters to estimate the year-wise population size of FSW, clients and the general population from 1995 to 2015¹⁷ (see supplementary table 1, text on pages 2–3, available online only). It is assumed that the FSW and clients are in the age group of 15–45 years and 15–50 years, respectively; and this cycle continues each year with the addition of a 15-year cohort and attrition of a 45 (50 for client)-year-old cohort. General all-cause mortality and HIV-specific mortality from existing Indian studies were applied to each of the population groups to estimate the annual attrition in numbers.^{18–19} The general population has been divided into married and unmarried cohorts using behavioural sentinel survey data (see supplementary table 1, available online only).^{20–21}

The probability of HIV transmission from men to women and vice versa has been estimated using the Weinstein equation, based on the Bernoulli theorem²² (see supplementary material, pages 5–8, available online only). This equation incorporates the effect of baseline HIV prevalence in the partner group, the risk of HIV transmission per unprotected sex act, condom use rate, effectiveness of condom, average number of sex partners and average number of sex acts per partner. Transmission coefficients for HIV per sex act are derived from epidemiological studies.^{23–27} We have assumed a rather wide uncertainty range for transmission coefficients as there is a lack of reliable data from India on the risk of HIV transmission per sex act. STI as a cofactor increases the risk of HIV transmission three times.^{25–28} This has been assumed to be one for a generic sexually transmitted disease (STD). This has also been used in other Indian studies.^{11–29} It was assumed that it would fit to the overall STD prevalence in different population groups, ie, FSW, clients and the general population.^{20–21–30–34}

Intervention effect

The targeted interventions result in an increase in safer sexual behaviour in terms of a reduction in multipartner sex and the greater use of condoms in commercial sex encounters. The

programme incorporates regular quarterly check-ups for detecting any STI. Presumptive and specific treatments for STI are given based on a specified protocol. This results in a reduction of STI prevalence among the FSW. We assume the effect of targeted interventions on condom use, STI treatment/prevalence, number of sex acts and number of sex partners of FSW.¹ Data from the two national rounds of behavioural surveillance (2001 and 2006), integrated biological and behavioural surveillance (IBBA) and demographic and health survey data have been used to compute the change in behavioural parameters (table 1).^{20–21–30–32–34}

The annual STI prevalence decline was estimated using programme FSW coverage, coverage of FSW for STI treatment and effectiveness of treatment. The probability of FSW getting re-infected with STI after being successfully treated for an episode is also incorporated in the model. FSW targeted intervention projects also treat partners of FSW.¹ However, as not all partners (clients) of FSW are treated, we assume that there would be a reduction in client STI prevalence, which is one third of FSW STI prevalence reduction. We assume that any reduction in STI prevalence in the general population women is not the direct effect of FSW targeted intervention projects and thus has not been modelled.

The effectiveness of the mass media in enhancing condom use has been derived from the literature.^{8–57} Both scenarios of FSW targeted intervention and mass media alone were assumed to operate in a setting in which ART was available. Coverage of ART was estimated from the literature (2% in 2002; 34% in 2008) and trends computed for years in which the coverage estimates were not available.^{56–58} ART has a lowering effect on HIV transmission as a result of decreased viral load and improved CD4 cell counts. Based on the observed reduction in viral load in blood plasma, experts estimate that ART reduces infectiousness by a factor of 2 to 8.^{47–59} Because the effect of ART on infectiousness is varied, our analysis uses the base assumption of a decrease in transmissibility by a factor of 4.

Costing

Cost data from an economic perspective on different interventions for all the intervening years is very scarce. Therefore,

Figure 1 Model structure for transmission of HIV epidemic from female sex workers to clients and general population, India. STD, sexually transmitted disease.




Table 1 Demographic, behavioural and biological parameters for the model to assess cost effectiveness of FSW targeted interventions in India

Parameter	Base value	Range		Source
		Lower	Upper	
Intervention parameters				
Efficacy of condom use				35–37
Vaginal	0.8	0.75	0.85	
Anal	0.7	0.65	0.75	
Oral	0.9	0.85	0.95	
Annual increase in condom use among FSW	0.091	0.073	0.099	21
Enrolment of FSW in targeted intervention projects (1995)	0.02			2
Enrolment of FSW in targeted intervention projects (2008)	0.53			38
Coverage of enrolled FSW for STI treatment (1995)	0.262			2
Effectiveness of STI treatment	0.85	0.68	0.98	39
Annual decline in risk behaviour				20 21 32
N (FSW)	0.02			
M (FSW)	0.03			
Behavioural parameters				
Proportion of female population involved in commercial sex	0.0056	0.039	0.006	21
Proportion of male population				
15–50 years sexually active	0.105	0.09	0.12	21
15–50 years having non-regular partner who reported commercial sex in 1 year	0.034	0.026	0.042	
Type of FSW-client sex acts				40 41
Vaginal	0.9	0.85	0.992	
Anal	0.05	0.003	0.08	
Oral	0.05	0.011	0.1	
Proportion condom use (1995)				
FSW	0.23	0.19	0.32	14
Married GP	0.022	0.009	0.06	30 33 34
Unmarried GP	0.014	0.01	0.018	
Annual change in condom use among married GP	0.09	0.05	0.14	30 34
Mean sex acts per FSW per client per year	3	2.25	3.75	Calculation based on IBBA, BSS-1, 2
Mean no of client per FSW per year (1995)	205	179	226	Calculation based on IBBA, Dandona et al ²⁹
Mean sex acts per client per FSW per year (1995)	2.81	2.5	3.0	Calculation based on IBBA, Dandona et al ²⁹
Mean no of FSW per client per year	16			Calculation based on IBBA, BSS-1, 2
Mean sex acts per partner per married GP per year	45			42
Mean sex acts per partner per unmarried GP per year	10			43
Mean no. of partners per married GP per year	1.019			31
Mean no. of partners per unmarried GP per year	0.046			31
Proportion of GP married-client sex act type				31
Vaginal	0.89			
Anal	0.009			
Oral	0.094			
Proportion of GP unmarried-client sex act type				31
Vaginal	0.89			
Anal	0.009			
Oral	0.094			
Biological parameters				
Proportion of FSW with				44 45
HIV	0.097	0.05	0.46	45
STI	0.42	0.35	0.47	46
Odds for STI among HIV infected	0.8			
Proportion of adolescent females with STI	0.041	0.004	0.162	31
Proportion of clients with				
HIV	0.047			Author analysis using HSS data
STI	0.26			21
Odds for STI among HIV infected	0.8			46

Continued

Table 1 Continued

Parameter	Base value	Range		Source
		Lower	Upper	
Probability of HIV transmission:				Range according to authors' estimates ^{25 27}
Male to female				
Vaginal	0.0014	0.0007	0.0021	
Anal	0.01	0.005	0.015	
Oral	0.0004	0.0002	0.0005	
Female to male				
Vaginal	0.0007	0.00035	0.0015	
Anal	0.001	0.0005	0.0015	
STI cofactor multiplicative effect	3	2	5	25 28
Probability of STI transmission:				47–50
Male to female				
Vaginal	0.25	0.2	0.5	
Anal	0.4	0.3	0.5	
Oral	0.1	0.05	0.15	
Female to male				
Vaginal	0.25			
Anal	0.4			
Proportion of female population (15–45 years) with				
HIV	0.002	0.2	0.3	Author analysis based on HSS data ⁴⁴
STI	0.2	0.3	0.5	
Cost parameters				
Total NACP budget				1
NACP-1 (1995–6)	80 crore			
NACP-2	1425 crore			
NACP-3	11583 crore			
Proportion of budget to mass media				1
No FSW intervention scenario				
NACP-1 (1995–6)	0.45			
NACP-2	0.25			
NACP-3	0.25			
FSW intervention scenario				
NACP-1 (1995–6)	0.45			
NACP-2	0.25			
NACP-3	0.1			
Average cost per FSW enrolled under targeted intervention				2 11 12 29 51–54(Bill and Melinda Gates Foundation and KHPT. Personal communication, 2010 and NACO. Personal communication, 2009)
NACP-1	794	773	1606	
NACP-2	1208	1612	3419	
NACP-3	2559			
Average cost per beneficiary for ART treatment				55
1995	15 168			
2008	9806			
Coverage of ART treatment				55 56
1995	0.01			
2008	0.34			

ART, antiretroviral therapy; BSS, behavioural sentinel surveillance; FSW, female sex worker; GP, general population; HSS, HIV sentinel surveillance; IBBA, integrated biological and behavioural surveillance; KHPT, Karnataka Health Promotion Trust; M (FSW) is the mean number of clients per FSW; N (FSW) is the mean sex acts per client per FSW per year; NACO, National AIDS Control Organization; NACP, National AIDS Control Programme; SRS, sample registration survey; STI, sexually transmitted infection.

we have estimated costs from a financial perspective. Cost in a no-targeted intervention scenario relates to the cost of mass media interventions and ART treatment for HIV patients. The targeted intervention arm additionally includes costs for implementing targeted interventions (see supplementary table 2, available online only). The cost of targeted interventions from 2003 onwards is a weighted average cost per FSW enrolled for both NACO and non-NACO including AVAHAN projects, using unit cost guidelines from NACO and AVAHAN (Bill and Melinda Gates Foundation and the Karnataka Health Promotion Trust (KHPT), personal communication, 2010 and A. Bhatia, personal communication, 2009). The weighting was done by the FSW population enrolled in each project according to the funding source. Cost data were validated with estimates on unit

costing from other studies.^{2 11 12 51} The cost of mass media interventions in the targeted intervention scenario was used directly from the NACO financial reports and programme implementation plan outlays published in annual reports¹ (table 1).

The model was parameterised with the cost of ART per beneficiary derived from a multicentric study from India (see supplementary figure 1, available online only).⁵⁵ The unit cost for ART was revised for recent years based on revised estimates of ART costing from NACO guidelines, programme implementation plans and the recent proposal submitted to the global fund.^{1 60}

All costs were converted to constant 2008 prices based on the consumer price index.⁶¹ The conversion factor for INR/US\$ has been used for 2009. Both costs and benefits have been discounted at 3%.

Valuing benefits

The primary endpoint for estimating benefits was a reduction in HIV cases. As a secondary endpoint, we used standard methods to estimate DALY averted resulting from FSW targeted interventions (see supplementary material, page 9, available online only).⁶²

Sensitivity analysis

Univariate sensitivity analysis was done to ascertain the influence of uncertainty in individual parameters on the summary measure of the incremental cost-effectiveness ratio (ICER). Key demographics (number of FSW, clients and general population), behavioural (condom use, sex behaviour), epidemiological (HIV prevalence among FSW, HIV and STD transmission probabilities, STD cofactor effect), intervention parameters (unit cost of targeted interventions, effectiveness of STI treatment) and discount rate (3% to 8%) were considered in sensitivity analysis.

Using MS Excel and Visual Basic, probabilistic sensitivity analysis was performed. We performed 1000 simulations using a uniform distribution of all parameters for which a plausible range was available (table 1 and supplementary table 1, available online only). Results from simulations were plotted on a cost-effectiveness plane. A cost-effectiveness acceptability curve was constructed to ascertain the probability of the targeted intervention being cost effective at varying threshold willingness-to-pay levels.

RESULTS

Targeted interventions for FSW result in a reduction of 47% (1.6 million) prevalent and 36% (2.7 million) cumulative HIV cases, respectively, in 2015. The prevalent and cumulative number of HIV cases in 2015, with targeted interventions for FSW, would be 1.8 and 4.7 million, respectively. Similarly, we found that FSW heterosexually acquired adult HIV prevalence in India would be 0.48% in 2015 (figure 2). The same prevalence would be reduced to 0.25% in the scenario of FSW targeted interventions. A reduction in non-condom use alone led to 32% and 40% declines in cumulative and prevalent numbers of HIV infections, respectively, in 2015.

The cumulative cost of implementing the FSW targeted interventions scenario from 1995 to 2015 was found to be INR61 693.5 million (US\$1991 million) (table 2) with an incremental cost of INR12 841 million (US\$285.3 million). From a health system perspective, the government and development




Figure 2 HIV prevalence in India, without targeted interventions (only mass media) and with targeted interventions for female sex workers along with mass media, 1995–2015. ANC, antenatal clinic; FSW, female sex worker; TI, targeted intervention.

Table 2 Cost effectiveness of FSW targeted interventions for HIV in India

Characteristic	Mass media	Mass media plus targeted intervention
Costs (million), (1995–2015)		
US\$	1085.6	1371
INR	48852.7	61693.5
Outcome measure (million), (2015)		
Cumulative HIV infections	7.4	4.7
DALY	91.7	65.1
Incremental cost effectiveness ratio		
Cost/HIV infection averted	US\$104 (INR4680)	
Cost/DALY averted	US\$10.7 (INR483)	
Incremental cost effectiveness ratio (discounted)		
Cost/HIV infection averted	US\$105.5 (INR4748)	
Cost/DALY averted	US\$10.9 (INR490)	

DALY, disability-adjusted life-year; FSW, female sex worker.

partners spend an average US\$104 (INR4680) per HIV infection averted and US\$10.7 (INR483) per DALY averted. Discounting at 3%, FSW targeted interventions cost US\$105.5 (INR4748) and US\$10.9 (INR490) per HIV case and DALY averted, respectively (table 2).

The risk of HIV (4.4–44.7) and STD (4.1–16.7) transmission per sex act, STD cofactor effect (4.4–25.2), client size (5.8–23.4), per unit cost of implementing targeted interventions (4.1–16.6) and FSW condom use (9.9–11.3) were the six parameters that induced maximum variation in the ICER per DALY averted (see supplementary figures 2 and 3, available online only).

Other factors such as HIV prevalence (FSW), the prevalence of sexually active men who report commercial sex activity, the type of sex act, other demographic parameters, efficacy of condom use and STI treatment, condom use among the general population and discount rate were found to have minimal impact on the overall cost effectiveness of FSW targeted interventions.


Targeted interventions for FSW are either dominant over the mass media-alone intervention (ie, cost saving) or cost effective at the current gross domestic product (GDP) level. Using 1000 simulations to account for all the parameters' uncertainty, there is nearly 70% probability for the targeted interventions to be cost effective at a willingness-to-pay threshold that is equal to the per capita GDP of India, ie, INR25 393 (figure 3).

DISCUSSION

We used a compartmental mathematical model to demonstrate the impact of preventive targeted interventions for FSW on HIV/AIDS in India and their cost effectiveness. Targeted interventions for FSW were found to be very cost effective from a health system perspective at US\$104 (INR4680) and US\$10.7 (INR483) per HIV case averted and per DALY averted, respectively. According to WHO guidelines on cost effectiveness, an intervention is considered to be cost effective if the value of ICER falls between 1 and 3 times the GDP per capita, and very cost effective if it is less than 1 GDP per capita. At a GDP per capita of INR25 393 (US\$564) for India, FSW targeted intervention is thus a very cost-effective strategy for HIV prevention.⁶³ Fung *et al*¹¹ also found similar findings in a model that was parameterised with data from Ahmedabad.

Our model made substantial improvements over the previous models for demonstrating the course of the HIV epidemic in India with or without interventions. Cost-effectiveness studies for preventive targeted interventions in India have used focal non-governmental organisation projects¹¹ or defined

Figure 3 Probability of targeted interventions to be cost effective at varying willingness-to-pay levels.



geographical area with characteristic behavioural and epidemiological parameters.²⁹ Our model is parameterised with India-specific demographic, behavioural and epidemiological data. The present model also accounts for the evolving nature of targeted interventions in terms of coverage and service provision. We modelled the effect of ART on viral load and its effect on the transmission of HIV, which is an improvement over some earlier models.⁶⁴ Unlike some previous models⁶⁴ the cost of ART in our study includes the cost for treating opportunistic infections, besides ART drugs, CD4 cell count testing, healthcare staff and other consumables.

Some of the earlier models have projected the HIV prevalence in India, which peaks to very high levels of almost 25 million cases.^{65–66} These observations could have resulted from overstated FSW numbers, or other behavioural and epidemiological parameters borrowed from African or other developed country settings. However, we used estimates on the FSW population based on revised mapping studies.³⁸ The sex behaviours of FSW and their clients were reviewed and calculated using data from national-level behavioural surveys and IBBA survey. Our per contact risk of transmission is derived from a review of the literature that also matches with assumptions used in other recent models of HIV transmission in India.^{6–11, 29} The estimates of HIV prevalence in our study are within the range of estimates from HIV sentinel surveillance in India. Our model shows 2.24 million HIV cases at the end of 2007 as against 2.31 million (1.8–2.9) according to revised estimates of HIV sentinel surveillance in India. We did not consider the HIV cases arising as a result of homosexual activity, parenteral and vertical transmission, or heterosexual networks of IDU and MSM. Our estimates should thus be viewed within this limitation. Adjusting for this limitation, the number of HIV-positive individuals in India would be in the range of 2.6–2.8 million assuming that the remaining modes of transmission constitute 25% to 15% of cases in India, respectively.¹

Our estimates of the cost effectiveness of FSW targeted interventions (US\$104 per HIV case averted) are similar to those of Fung *et al*¹¹ (US\$34–133) from Gujarat and Creese *et al*⁶⁴ (US \$79–160) from Cameroon but different from those reported by Hogan *et al*¹⁰ for south east Asian (SEAR D) countries, which found targeted interventions to cost \$45, \$47 and \$50 international

(2000) at coverage level of 50%, 80% and 95%, respectively. In terms of international dollars, we found FSW targeted interventions to cost \$244 international (2005) per HIV infection averted.⁶⁷ However, these differences are based on differing methodology and assumptions of two studies. Hogan *et al*¹⁰ have hypothetically assumed a scenario of ‘no condom use’ and ‘no STI treatment’ to compare with an alternative scenario of targeted interventions for FSW.

Certain limitations of the present research need to be highlighted. A wide range of STI cofactor effect on HIV transmission as a result of different bacterial and viral STI has been noted.²⁸ However, we have assumed a constant STI cofactor multiplicative effect of 3 for a generic STI. A higher viral load enhances HIV transmission during the sex act.⁴⁷ Our model uses average per sex act transmission probabilities for HIV acquisition among uninfected individuals. However, we do factor the viral-lowering effect of ART treatment and thus lowered transmission of HIV among those on treatment.

Our predicted HIV prevalence among FSW and clients is higher than those from HIV sentinel surveillance and those

Key messages

- ▶ We assessed the cost effectiveness of targeted interventions for FSW in India, which is a strategy for the prevention of the focused HIV epidemic among high-risk groups through peer-led behaviour change communication.
- ▶ Our study found that the targeted interventions for FSW reduced the prevalence of HIV infections in India by 47% from 1995 to 2015.
- ▶ Using a deterministic model, we found that targeted intervention for FSW is a very cost-effective strategy for HIV prevention in India, with an incremental cost of US\$10.7 (INR483) per DALY averted.
- ▶ Under all uncertainties involved with parameter estimation, we found a 90% probability of targeted interventions for FSW to be cost effective at a willingness-to-pay threshold equal to 1.5 times per capita GDP of India.

reported by Alary *et al*¹⁵ using the IBBA data. FSW HIV prevalence in increasing order is found from HIV sentinel surveillance followed by Alary *et al*¹⁵ (IBBA surveillance data) and is highest according to our study. We would also like to emphasise that the antenatal clinic HIV prevalence predicted from our model matches those from the HIV sentinel surveillance data and the IBBA results. This could be due to the lesser representative nature of sentinel sites for high-risk groups in HIV sentinel surveillance, especially during early years; and the non-random nature of sample collection (for further discussion, refer to the supplementary material available online only).

The present study was limited to an analysis of the cost effectiveness of FSW targeted interventions. Two more approaches for future research on the resource allocation of India's HIV prevention programme are suggested. First would be to look at a broader level and ascertain the impact and cost effectiveness of different preventive interventions such as targeted interventions for MSM, IDU, voluntary counselling and testing centres, mass media, etc. This will inform a better resource allocation decision making in India's National AIDS Control Programme. Second, similar interventions (eg, targeted interventions for FSW) are delivered with different intensity and different strategy (such as NACO-funded projects and AVAHAN-funded projects). Therefore, it is suggested to ascertain the impact and cost effectiveness of different approaches of service delivery at the micro level.

Funding This study was funded by the National AIDS Control Organization, New Delhi, India.

Competing interests All authors have received support from the National AIDS Control Organization, New Delhi, India, for the submitted work.

Ethics approval This study was conducted with the approval of the Ethics Committee, PGIMER, Chandigarh, India.

Contributors S. Prinja: Developed the model, interpreted the results and wrote the manuscript. PB: Contributed to development of the model, reviewed literature and performed sensitivity analysis. RK, SMM, MK, S.Panda and IG: Contributed to conceptualisation of the model, results interpretation, reviewed and revised the manuscript. SR and SC: Reviewed literature for estimating parameters and reviewed the manuscript. All authors approved the final version of manuscript being submitted to the journal for publication.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- NACO. *Program Implementation Plan National AIDS Control Program III*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, 2006.
- Kumar R, Mehendale SM, Panda S. *Impact assessment of targeted interventions on HIV prevention in India: Project Report-Part 1*. Chandigarh: PGIMER School of Public Health, 2010.
- Reza-Paul S, Beattie T, Syed HU, *et al*. Declines in risk behaviour and sexually transmitted infection prevalence following a community-led HIV preventive intervention among female sex workers in Mysore, India. *AIDS* 2008;**22**(Suppl 5):S91–100.
- Basu I, Jana S, Rotheram-Borus MJ, *et al*. HIV prevention among sex workers in India. *J Acquir Immune Defic Syndr* 2004;**36**:845–52.
- Bhave G, Lindan C, Hudes E, *et al*. Impact of an intervention on HIV, sexually transmitted diseases and condom use rates among sex workers in Bombay, India. *AIDS* 1995;**9**(Suppl 1):S21–30.
- Boily MC, Pickles M, Vickerman P, *et al*. Using mathematical modelling to investigate the plausibility of attributing observed antenatal clinic declines to a female sex worker intervention in Karnataka state, India. *AIDS* 2008;**22**(Suppl 5):S149–64.
- Nagelkerke NJ, Jha P, de Vlas SJ, *et al*. Modelling HIV/AIDS epidemics in Botswana and India: impact of interventions to prevent transmission. *Bull WHO* 2002;**80**:89–96.
- Bollinger L, Cooper-Arnold K, Stover J. Where are the gaps? The effects of HIV-prevention interventions on behavioral change. *Stud Fam Plann* 2004;**35**:27–38.
- Walker D. Cost and cost-effectiveness of HIV/AIDS prevention strategies in developing countries: is there an evidence base? *Health Policy Plan* 2003;**18**:4–17.
- Hogan DR, Baltussen R, Hayashi C, *et al*. Cost effectiveness analysis of strategies to combat HIV/AIDS in developing countries. *BMJ* 2005;**331**:1431–7.
- Fung IC, Guinness L, Vickerman P, *et al*. Modelling the impact and cost-effectiveness of the HIV intervention programme amongst commercial sex workers in Ahmedabad, Gujarat, India. *BMC Public Health* 2007;**7**:195.
- Dandona L, Kumar SP, Ramesh Y, *et al*. Changing cost of HIV interventions in the context of scaling-up in India. *AIDS* 2008;**22**(Suppl 1):S43–9.
- NACO. *Annual Report 2009–10*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, 2010.
- Venkataramana CB, Sarada PV. Extent and speed of spread of HIV infection in India through the commercial sex networks: a perspective. *Trop Med Int Health* 2001;**6**:1040–61.
- Alary M, Jayachandran AA, Lowndes CM, *et al*. Ecological analysis of the association between high-risk population parameters and HIV prevalence among pregnant women enrolled in sentinel surveillance in four Southern India states. *Sex Transm Infect* 2010;**86**(Suppl 1):i10–16.
- WHO. *Making choices in health: WHO guide to cost-effectiveness analysis*. Geneva: World Health Organization, 2003.
- RGI. *Population projections for India and States 2001–2026 (revised December 2006)*. New Delhi: Registrar General and Census Commissioner India, 2006.
- Jha P, Kumar R, Khera A, *et al*. HIV mortality and infection in India: estimates from nationally representative mortality survey of 1.1 million homes. *BMJ* 2010;**340**:c621.
- SRS. *Estimated age-specific death rates by sex in India and Southern States 1994*. Registrar General of India Sample Registration System. http://www.kerala.gov.in/statistical/vitalstatistics/w_1.15.pdf (accessed 3 Jan 2010).
- NACO. *National Behavioural Surveillance Survey (2001)*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, 2001.
- NACO. *National Behavioural Surveillance Survey (BSS) 2006: female sex workers and their clients*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, 2006.
- Weinstein M, Graham J, Siegel J, *et al*. Cost-effectiveness analysis of AIDS prevention programs: concepts, complications and illustrations. In: Turner C, Miller H, Moses L, eds. *AIDS: sexual behaviour and intravenous drug use*. Washington, DC: National Academy Press, 1989.
- Boily MC, Baggaley RF, Wang L, *et al*. Heterosexual risk of HIV-1 infection per sexual act: systematic review and meta-analysis of observational studies. *Lancet Infect Dis* 2009;**9**:118–29.
- Baggaley RF, White RG, Boily MC. HIV transmission risk through anal intercourse: systematic review, meta-analysis and implications for HIV prevention. *Int J Epidemiol* 2010;**39**:1048–63.
- Gouws E, White P, Stover J, *et al*. Short term estimates of adult HIV incidence by mode of transmission: Kenya and Thailand as examples. *Sex Transm Infect* 2006;**82**(Suppl 3):iii51–5.
- Vittinghoff E, Douglas J, Judson F, *et al*. Per-contact risk of human immunodeficiency virus transmission between male sexual partners. *Am J Epidemiol* 1999;**150**:306–11.
- Royce RA, Sena A, Cates W Jr, *et al*. Sexual transmission of HIV. *N Engl J Med* 1997;**336**:1072–8.
- Korenromp EL, De Vlas SJ, Nagelkerke N, *et al*. Estimating the magnitude of STD cofactor effects on HIV transmission, how well can it be done? *Sex Transm Dis* 2001;**28**:613–21.
- Dandona L, Kumar S, Kumar G, *et al*. Economic analysis of HIV prevention interventions in Andhra Pradesh state of India to inform resource allocation. *AIDS* 2009;**23**:233–42.
- IIPS. *National Family Health Survey 2 (1998–99)*. Mumbai: International Institute of Population Sciences, and ORC Macro, 2000.
- NACO. *National Behavioural Surveillance Survey (BSS) 2006: General Population*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, Government of India, 2006.
- ICMR FHI. *National Interim Summary Report India, Integrated Behavioural and Biological Assessment Round 1 (2005–2007)*. New Delhi: Indian Council of Medical Research New Delhi and Family Health International, 2007.
- IIPS. *National Family Health Survey 1 (1992–93)*. Mumbai: International Institute of Population Sciences, and ORC Macro, 1994.
- IIPS. *National Family Health Survey 3 (2005–06)*. Mumbai: International Institution of Population Sciences and ORC Macro, 2007.
- Weller S, Davis K. Condom effectiveness in reducing heterosexual HIV transmission. *Cochrane Database Syst Rev* 2002;(1):CD003255.
- Pinkerton SD, Abramson PR. Effectiveness of condoms in preventing HIV transmission. *Soc Sci Med* 1997;**44**:1303–12.
- Dandona L, Sisodia P, Kumar SG, *et al*. HIV prevention programmes for female sex workers in Andhra Pradesh, India: outputs, cost and efficiency. *BMC Public Health* 2005;**5**:98.
- NACO. *HIV Sentinel Surveillance and HIV Estimation in India 2007 A Technical Brief*. New Delhi: National AIDS Control Organization, Ministry of Health and Family Welfare, 2008. http://www.naonline.org/upload/Publication/MGE%20Surveillance,%20Research/HIV%20Sentinel%20Surveillance%20and%20HIV%20Estimation%202007_A%20Technical%20Brief.pdf (accessed 15 May 2010).
- Pettifor A, Walsh J, Wilkins V, *et al*. How effective is syndromic management of STDs? a review of current studies. *Sex Transm Dis* 2000;**27**:371–85.
- Matheou A. *A blind spot in HIV prevention – female anal sex*. New Delhi: India HIV/AIDS Alliance, 2010.
- Bloem M, Hoquea E, Khanama L, *et al*. HIV/AIDS and female street based sex workers in Dhaka city: what about their clients? In: Caldwell J, Caldwell P, Anarfi J, *et al*, eds. *Resistances to behavioural change to reduce HIV/AIDS Infection*. Canberra. Health Transition Centre, National Centre for Epidemiology and Population Health, Australian National University, 1999:197–210.

42. **Survey GS.** *Durex Global Sex Survey 2005.* <http://www.durex.com/cm/gss2005result.pdf> (accessed 30 Mar 2010).
43. **Jejeebhoy SJ.** Adolescent sexual and reproductive behavior: a review of the evidence from India. *Soc Sci Med* 1998;**46**:1275–90.
44. **NICED.** *Annual Report.* Kolkata: National Institute of Cholera and Enteric Diseases and Indian Council of Medical Research, 1986–7.
45. **Kosambiya J,** Desai V, Thakor H, *et al.* Risk markers for presence of sexually transmitted infections in female sex workers, India. *Indian J Sex Transm Dis* 2005;**26**:26–32.
46. **Bollinger RC,** Brookmeyer RS, Mehendale SM, *et al.* Risk factors and clinical presentation of acute primary HIV infection in India. *JAMA* 1997;**278**:2085–9.
47. **Chakraborty H,** Sen PK, Helms RW, *et al.* Viral burden in genital secretions determines male-to-female sexual transmission of HIV-1: a probabilistic empiric model. *AIDS* 2001;**15**:621–7.
48. **Mann JR,** Stine CC, Vessey J. The role of disease-specific infectivity and number of disease exposures on long-term effectiveness of the latex condom. *Sex Transm Dis* 2002;**29**:344–9.
49. **Kretzschmar M,** Welte R, van den Hoek A, *et al.* Comparative model-based analysis of screening programs for *Chlamydia trachomatis* infections. *Am J Epidemiol* 2001;**153**:90–101.
50. **Quinn TC,** Gaydos C, Shepherd M, *et al.* Epidemiologic and microbiologic correlates of *Chlamydia trachomatis* infection in sexual partnerships. *JAMA* 1996;**276**:1737–42.
51. **Bhat R,** Sudhakar K, Thomas K. Analysis of programme targets and expenditures. *Mid-Term Review Meeting for National AIDS Control Program III; 2010.* New Delhi, 16 November 2009.
52. **Guinness L,** Kumaranayake L, Hanson K. A cost function for HIV prevention services: is there a 'u'-shape? *Cost Effic Resour Alloc* 2007;**5**:13.
53. **Guinness L,** Kumaranayake L, Rajaraman B, *et al.* Does scale matter? The costs of HIV-prevention interventions for commercial sex workers in India. *Bull WHO* 2005;**83**:747–55.
54. **Chandrashekar S,** Guinness L, Kumaranayake L, *et al.* The effects of scale on the costs of targeted HIV prevention interventions among female and male sex workers, men who have sex with men and transgenders in India. *Sex Transm Infect* 2010;**86** (Suppl 1):i89–94.
55. **Gupta I,** Trivedi M, Kandamathan S. *An analysis of recurrent costs of the Free ART Program of the Government of India.* New Delhi: Health Policy Research Unit, Institute of Economic Growth, 2006.
56. **WHO.** *HIV/AIDS in South East Asia Region.* New Delhi: World Health Organization Regional Office for South East Asia Region, 2009.
57. **Goldstein S,** Scheepers E. Soul City 4 impact evaluation. *AIDS.* <http://www.soulcity.org.za> (accessed 23 Dec 2010).
58. **Gupta I,** Sankar D. *Treatment seeking behaviour and the willingness to pay for antiretroviral therapy in India.* New Delhi: Institute of Economic Growth Working Paper, 2003.
59. **Over M,** Marseille E, Sudhakar K, *et al.* Antiretroviral therapy and HIV prevention in India: modeling costs and consequences of policy options. *Sex Transm Dis* 2006;**33** (Suppl 10):S145–52.
60. **NACO.** *Proposal submitted to Global Fund, Round 4 RCC.* New Delhi: National AIDS Control Organization, 2006.
61. **RBI.** *Handbook of Statistics on Indian Economy.* New Delhi: Reserve Bank of India. http://rbidocs.rbi.org.in/rdocs/Publications/PDFs/OHANDB210910_F.pdf (accessed 15 May 2010).
62. **Fox-Rushby JA,** Hanson K. Calculating and presenting disability adjusted life years (DALYs) in cost-effectiveness analysis. *Health Policy Plan* 2001;**16**:326–31.
63. **WHO.** *Tough choices: investing in health for development experiences from the national follow-up to the macroeconomic commission on health.* Geneva, Switzerland: WHO, 2006.
64. **Creese A,** Floyd K, Alban A, *et al.* Cost-effectiveness of HIV/AIDS interventions in Africa: a systematic review of the evidence. *Lancet* 2002;**359**:1635–43.
65. **Rao C,** Srivenkataramana T. Projection of HIV infections in India: an alternative to back-calculation. *Curr Sci* 2001;**81**:1302–7.
66. **Jha P.** *Project Appraisal Document: Second National HIV/AIDS Control Project.* New Delhi: World Bank, 1999.
67. **WHO.** *Purchasing Power Parity 2005.* 2005. <http://www.who.int/choice/costs/ppp/en/> (accessed 28 Apr 2010).



Cost effectiveness of targeted HIV prevention interventions for female sex workers in India

Shankar Prinja, Pankaj Bahuguna, Shalini Rudra, et al.

Sex Transm Infect published online March 29, 2011
doi: 10.1136/sti.2010.047829

Updated information and services can be found at:
<http://sti.bmj.com/content/early/2011/03/29/sti.2010.047829.full.html>

	<i>These include:</i>
Data Supplement	"Web Only Data" http://sti.bmj.com/content/suppl/2011/03/28/sti.2010.047829.DC1.html
References	This article cites 37 articles, 12 of which can be accessed free at: http://sti.bmj.com/content/early/2011/03/29/sti.2010.047829.full.html#ref-list-1
P<P	Published online March 29, 2011 in advance of the print journal.
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>